

Section 3.10
Related Rates
2 Lecture

Dr. Abdulla Eid

College of Science

MATHS 101: Calculus I

Rate of change

Recall:

- If a **line** has a slope $m = 3$, it means that for every one step to the right, we move along the line 3 steps up. In this case, as x **increases**, y **increases**.
- If a **line** has a slope $m = -2$, it means that for every 1 step to the right, we move along the line 2 steps down. In this case, as x **increases**, y **decreases**.

Exercise 1

If a line has a slope $\frac{2}{-3}$. What does that tell you about the line?

For general function $y = f(x)$, for every step to the right, how many steps do we go up/down? How do we measure that change in y ?

If x changes by 1, an estimate of the change in y is $\frac{dy}{dx}$.

Definition 2

The derivative of $y = f(x)$ can be interpreted as *rate of change* of y in terms of x .

In this section, we have our variables are functions in t . so for example, we have

- 1 $\frac{dx}{dt}$ is the rate change in x with respect to the time t . If $\frac{dx}{dt} = 4$, it means that for every one unit in time, x increases by 4 units.
- 2 $\frac{dr}{dt}$ is the rate change in r with respect to the time t . If $\frac{dr}{dt} = -2$ cm/second, it means that for every one second, r **decreases** 2 cm.

Example 3

If $y = x^2$ and $\frac{dx}{dt} = 3$. What is $\frac{dy}{dt}$ when $x = -1$?

Solution:

Given: $\frac{dx}{dt} = 3$ and $x = -1$.

Required: $\frac{dy}{dt}$.

Relation: $y = x^2$

$$\frac{dy}{dt} = 2x \frac{dx}{dt}$$

$$\frac{dy}{dt} = 2(-1)(3) = -6$$

Exercise 4

If $z = 3y^3$ and $\frac{dz}{dt} = 10$. What is $\frac{dy}{dt}$ when $y = 2$?

Example 5

Air is being pumped in a spherical balloon so that its volume increases at a rate of $500 \text{ cm}^3/\text{second}$. How fast is the radius of the balloon increases when the diameter is 50 cm ? (Note that the volume of the sphere is $V = \frac{4}{3}\pi r^3$).

Solution:

Given: $\frac{dV}{dt} = 500$ and $D = 50 \rightarrow r = \frac{D}{2} = 25$.

Required: $\frac{dr}{dt}$.

Relation: $V = \frac{4}{3}\pi r^3$

$$\frac{dV}{dt} = \frac{4}{3}\pi(3r^2) \frac{dr}{dt}$$

$$500 = 4\pi(25)^2 \frac{dr}{dt} \rightarrow \frac{dr}{dt} = \frac{500}{4\pi(25)^2} \text{ cm / second}$$

Exercise 6

A circular disk is expanding at a rate $10 \text{ cm}^2/\text{second}$. Find the rate of change of the Diameter of the disk when the radius is 5 cm.

Dr. Abdulla Eid

Example 7

Each side of a square is increasing at a rate of 6 cm/second. At what rate is the area of the square increases when the area of the square is 16 cm².

Solution:

Given: $\frac{dx}{dt} = 6$ and $A = 16$.

Required: $\frac{dA}{dt}$.

Relation: $A = x^2$

$$\begin{aligned}\frac{dA}{dt} &= 2x \frac{dx}{dt} \\ \frac{dA}{dt} &= 2(4)(6) = \frac{dA}{dt} = 24 \text{ cm}^2 / \text{second}\end{aligned}$$

Exercise 8

The length ℓ of a rectangle is decreasing at the rate of 2 cm/second while the width w is increasing at the rate of 2 cm/second when $\ell = 12$ cm and $w = 5$ cm. Find the rate of change of

- 1 Area
- 2 Perimeter
- 3 Length of the diagonal

Dr. Abdulla

Example 9

A ladder is 5 m long rests against a vertical wall. If the bottom of the ladder moves away at rate 0.5 m/second. How fast the top of the ladder slides down when the bottom of the ladder is 4 m from the wall.

Solution:

Given: $\frac{dx}{dt} = 0.5$ and $l = 5$, $x = 4$.

Required: $\frac{dy}{dt}$.

Relation: $x^2 + y^2 = 25$

$$2x \frac{dx}{dt} + 2y \frac{dy}{dt} = 2(4)(0.5) + 2(3) \left(\frac{dy}{dt} \right) = 0$$

$$4 + 6 \frac{dy}{dt} = 0 \rightarrow \frac{dy}{dt} = \frac{-4}{6} \text{ m / second}$$

Example 10

An airplane is moving parallel to the ground on elevation 10 km with a rate of 6 km/minute. Find the rate of change for the angle where an observer observes the airplane when his angle is $\frac{\pi}{3}$.

Solution:

Given: $\frac{dx}{dt} = 6$ and $y = 10$, $\theta = \pi/3$.

Required: $\frac{d\theta}{dt}$.

Relation: $\tan \theta = \frac{10}{x}$ (hence $x = \frac{10}{\tan(\frac{\pi}{3})}$)

$$\sec^2 \theta \cdot \frac{d\theta}{dt} = \frac{-10}{x^2} \cdot \frac{dx}{dt}$$

$$2 \frac{d\theta}{dt} = \frac{-10}{(\)^2} \cdot 6 \rightarrow \frac{d\theta}{dt} = \text{radian / minute}$$

Exercise 11

The area of a triangle with sides a and b and contained angle θ is given by

$$A = \frac{1}{2}ab \sin \theta$$

If a increases at a rate of 2.5 cm/min, b increases at a rate of 1.5 cm/min, and θ increases at a rate of 0.2 rad/min. How fast is the area increasing when $a = 2$ cm, $b = 3$ cm, and $\theta = \frac{\pi}{3}$.

Dr. Abdu

Example 12

A cube's surface area increases at a rate of $72 \text{ in}^2/\text{second}$. At what rate the cube volume changes when the edge of the cube is 3 in ? (Note that the volume of the cube is $V = x^3$ and the surface area is $S = 6x^2$)

Solution:

Given: $\frac{dS}{dt} = 72$ and $x = 3$.

Required: $\frac{dV}{dt}$.

Relation: $V = x^3$

$$\frac{dV}{dt} = (3x^2) \frac{dx}{dt} \rightarrow \frac{dV}{dt} = 27 \frac{dx}{dt}$$

$$S = 6x^2 \rightarrow \frac{dS}{dt} = 12x \frac{dx}{dt} \rightarrow 72 = 36 \frac{dx}{dt} \rightarrow \frac{dx}{dt} = 2$$

$$\frac{dV}{dt} = 27(2) = 54 \text{ in}^3/\text{second}$$