

Section 5.4
Initial value problems
1 Lecture

Dr. Abdulla Eid

College of Science

MATHS 104: Mathematics for Business II

Differential equations

Definition 1

The *Initial value problem* is the problem of finding the function y given the derivative y' and *initial condition* $y(a) = b$.

Idea: Integrate y' and find the general antiderivative and then substitute $x = a$ and $y = b$ to find the function y .

Note: The initial value problem is subclass of a bigger problem in mathematics called solving **differential equations**.

Example 2

If y satisfies the given condition, find $y(x)$.

$$\frac{dy}{dx} = 6x^2 - 8x + 12 \text{ and } y(0) = 2$$

Solution: We integrate to find y .

$$y = \int (6x^2 - 8x + 12) dx$$

$$y = 2x^3 - 4x^2 + 12x + C$$

$$2 = 2(0)^3 - 4(0)^2 + 12(0) + C$$

$$2 = C$$

$$y = 2x^3 - 4x^2 + 12x + 2$$

Exercise 3

If y satisfies the given condition, find $y(x)$.

$$\frac{dy}{dx} = 9x^2 - 4x + 5 \text{ and } y(-1) = 0.$$

Dr. Abdulla Eid

Example 4

If y satisfies the given condition, find $y(x)$.

$$y'' = 6x + 4 \text{ and } y'(0) = 1, y(0) = 5$$

Solution: We integrate to find y .

$$y' = \int (6x + 4) dx$$

$$y' = 3x^2 + 4x + C$$

$$1 = 3(0)^2 + 4(0) + C \rightarrow C = 1$$

$$y' = 3x^2 + 4x + 1$$

$$y = \int (3x^2 + 4x + 1) dx$$

$$y = x^3 + 2x^2 + x + D$$

$$5 = y(0) \rightarrow D = 5$$

$$y = x^3 + 2x^2 + x + 5$$

Exercise 5

If y satisfies the given condition, find $y(t)$.

$$\frac{d^2y}{dt^2} = \cos t + \sin t \text{ and } y'(\pi) = 0, y(\pi) = 1$$

Dr. Abdulla Eid

Example 6

If y satisfies the given condition, find $y(x)$.

$$\frac{dy}{dx} = e^{-x} + \frac{1}{\sqrt{1-x^2}} \text{ and } y(0) = \pi$$

Solution: We integrate to find y .

$$y = \int \left(e^{-x} + \frac{1}{\sqrt{1-x^2}} \right) dx$$

$$y = -e^{-x} + \sin^{-1} x + C$$

$$\pi = -e^{-0} + \sin^{-1}(0) + C$$

$$\pi = -1 + 0 + C$$

$$\pi - 1 = C$$

$$y = -e^{-x} + \sin^{-1} x + \pi - 1$$

Exercise 7

If y satisfies the given condition, find $y(x)$.

$$y' = \frac{3}{1+x^2}, \quad y'(1) = 5$$

Dr. Abdulla Eid