

# Section 4.5

## Dimension

Dr. Abdulla Eid

College of Science

MATHS 211: Linear Algebra

## Goal:

- 1 Define the dimension of a vector space.
- 2 Some Important Theorem.

Dr. Abdulla Eid

# Basis

## Definition 1

Let  $V$  be a vector space. Let  $\mathcal{B} = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$  be a **basis**. The number  $n$  is called the **dimension** of the vector space  $V$ , and denoted by  $\dim(V)$ .

Note: This number  $n$  is independent of the chosen basis

## Theorem 2

*All bases for a finite-dimensional vector space have the same number of vectors.*

### Lemma 3

Let  $V$  be a vector space. Let  $\mathcal{B} = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$  be a basis.

- 1 If a set has more than  $n$  vectors, then it is linearly dependent.
- 2 If a set has fewer than  $n$  vectors, then it does not span  $V$ .

# Standard Basis

Note:

- The standard basis for  $\mathbb{R}^2$  is  $e_1, e_2$ , where

$$e_1 = (1, 0) \text{ and } e_2 = (0, 1)$$

So  $\dim(\mathbb{R}^2) = 2$ .

- The standard basis for  $\mathbb{R}^3$  is  $e_1, e_2, e_3$ , where

$$e_1 = (1, 0, 0), e_2 = (0, 1, 0) \text{ and } e_3 = (0, 0, 1)$$

$\dim(\mathbb{R}^3) = 3$ .

- The standard basis for  $\mathbb{R}^n$  is  $e_1, e_2, e_3, \dots, e_n$ , where

$$e_1 = (1, 0, \dots, 0), e_2 = (0, 1, \dots, 0), e_3 = (0, 0, 1, 0, \dots, 0) \text{ and } e_n = (0, 0, \dots, 1)$$

$\dim(\mathbb{R}^n) = n$ .

- The standard basis for  $\mathbb{P}_2$  is  $1, X, X^2$ .

$\dim(\mathbb{P}_2) = 3$ .

- The standard basis for  $\mathbb{P}_n$  is  $1, X, X^2, \dots, X^n$ .

$\dim(\mathbb{P}_n) = n + 1$ .

The standard basis for  $\text{Mat}(2, 2, \mathbb{R})$  is  
 $\dim(\text{Mat}(2, 2, \mathbb{R})) = 4$ . and in general  $\dim(\text{Mat}(m, n, \mathbb{R})) = mn$ .

Dr. Abdulla Eid

### Example 4

Find bases for the subspace of  $\mathbb{R}^3$  given by the plane  $2x + 4y - 3z = 0$

Dr. Abdulla Eid

## Homework 5

### Example 5

Find bases for the subspace of  $\mathbb{R}^3$  given by the plane  $x + z = 0$

Dr. Abdulla Eid



### Example 6

Find bases for the subspace of  $\mathbb{R}^3$  given by the plane  $x = 4t, y = 2t, z = -t$ .

Dr. Abdulla Eid

### Example 7

Find bases for the subspace of  $\mathbb{R}^3$  given by all vectors of the form  $(a, b, c)$ , where  $c = a - b$ .

Dr. Abdulla Eid

### Example 8

Find the dimension of the subspace  $W$  of  $\mathbb{R}^4$ , given by all vectors of the form  $(0, a, b, c)$ .

Dr. Abdulla Eid

### Example 9

Find the dimension of the subspace  $W$  of  $\mathbb{R}^4$ , given by all vectors of the form  $(a, b, c, d)$ , where  $d = a + 2b$ ,  $c = 3a - b$ .

Dr. Abdulla Eid

## Homework 5

### Example 10

Find the dimension of the subspace  $W$  of  $\mathbb{R}^4$ , given by all vectors of the form  $(a, b, c, d)$ , where  $c = a, b = d = -a$ .

Dr. Abdulla E

### Example 11

Find the dimension of the subspace  $W$  of  $\text{Mat}(n, n, \mathbb{R})$ , given by all diagonal matrices.

Dr. Abdulla Eid

### Example 12

Find the dimension of the subspace  $W$  of  $\text{Mat}(n, n, \mathbb{R})$ , given by all symmetric matrices.

Dr. Abdulla Eid

### Example 13

Find the dimension of the subspace  $W$  of  $\text{Mat}(n, n, \mathbb{R})$ , given by all upper triangular matrices.

Dr. Abdulla Eid



### Example 14

Find the dimension of the subspace  $W$  of  $\mathbb{P}_n$ , given by all polynomials with a horizontal tangent at  $x = 0$ .

Dr. Abdulla Eid

# Infinite Dimensional Spaces

## Example 15

Let  $V = \text{Maps}(\mathbb{R}, \mathbb{R})$ . Show that for every positive integer  $n$ , one can find  $n + 1$  independent functions.

Conclusion:  $\text{Maps}(\mathbb{R}, \mathbb{R})$  is infinite-dimensional. So do  $C^i(\mathbb{R}, \mathbb{R})$  and  $C^\infty(\mathbb{R}, \mathbb{R})$ .

# Plus/Minus Theorem

## Theorem 16

Let  $S$  be a nonempty set of vectors in a vector space  $V$ .

- 1 If  $S$  is a linearly independent set, and  $\mathbf{v} \in V$  that is outside  $\text{Span}(S)$ , then the set  $S \cup \{\mathbf{v}\}$  is still linearly independent.
- 2 If  $\mathbf{v}$  is a vector in  $S$  that is expressible as a linear combination of other vectors in  $S$ , then

$$\text{Span}(S) = \text{Span}(S - \{\mathbf{v}\})$$

### Example 17

Enlarge the set  $\left\{ \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 2 \\ -1 \\ 3 \end{pmatrix} \right\}$  to produce a basis for  $\mathbb{R}^3$ .

Dr. Abdulla Eid

## Homework 5

### Example 18

Enlarge the set  $\left\{ \begin{pmatrix} 5 \\ 3 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix} \right\}$  to produce a basis for  $\mathbb{R}^3$ .

Dr. Abdulla