

Section 6.1

Inner Product

Dr. Abdulla Eid

College of Science

MATHS 211: Linear Algebra

Goal:

- 1 Cauchy Schwarz Inequality.
- 2 Angle between vectors.
- 3 Properties of length and distance.
- 4 Orthogonality.
- 5 Orthogonal Complement.

Dr. Abdulla Eid

Cauchy Schwarz Inequality

Theorem 1

If \mathbf{u}, \mathbf{v} are two vectors, then

$$|\langle \mathbf{u}, \mathbf{v} \rangle| \leq \|\mathbf{u}\| \|\mathbf{v}\|$$

Proof: Let $a = \|\mathbf{u}\|^2$, $b = 2\langle \mathbf{u}, \mathbf{v} \rangle$, and $c = \langle \mathbf{v}, \mathbf{v} \rangle$. Consider $\langle t\mathbf{u} + \mathbf{v}, t\mathbf{u} + \mathbf{v} \rangle \geq 0$ so the discriminant is less than or equal to zero.

Dr. Abdulla Eid

Consequences of Cauchy Schwarz Inequality

$$|\langle \mathbf{u}, \mathbf{v} \rangle| \leq \|\mathbf{u}\| \|\mathbf{v}\|$$

$$\frac{|\langle \mathbf{u}, \mathbf{v} \rangle|}{\|\mathbf{u}\| \|\mathbf{v}\|} \leq 1$$

$$-1 \leq \frac{\langle \mathbf{u}, \mathbf{v} \rangle}{\|\mathbf{u}\| \|\mathbf{v}\|} \leq 1$$

$$\cos \theta = \frac{\langle \mathbf{u}, \mathbf{v} \rangle}{\|\mathbf{u}\| \|\mathbf{v}\|} \text{ and } 0 \leq \theta \leq \pi$$

Definition 2

The angle θ between \mathbf{u} and \mathbf{v} is defined by

$$\theta = \cos^{-1} \left(\frac{\langle \mathbf{u}, \mathbf{v} \rangle}{\|\mathbf{u}\| \|\mathbf{v}\|} \right)$$

Find the angle

Example 3

Find the cosine of the angle between \mathbf{u} and \mathbf{v} , with the standard inner product.

$$\textcircled{1} \quad \mathbf{u} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} \text{ and } \mathbf{v} = \begin{pmatrix} 4 \\ 4 \\ -4 \end{pmatrix}.$$

$$\textcircled{2} \quad \mathbf{u} = \begin{pmatrix} 0 \\ 1 \\ 2 \\ 3 \end{pmatrix} \text{ and } \mathbf{v} = \begin{pmatrix} 5 \\ 2 \\ -3 \\ -4 \end{pmatrix}.$$

$$\textcircled{3} \quad p = 1 + 3X - 5X^2, \quad q = 2 - 3X^2.$$

$$\textcircled{4} \quad U = \begin{pmatrix} 2 & 4 \\ 3 & 2 \end{pmatrix}, \quad V = \begin{pmatrix} 0 & 1 \\ 0 & -2 \end{pmatrix}$$

Properties of Length and Distance

Theorem 4

If $\mathbf{u}, \mathbf{v}, \mathbf{w}$ are three vectors, then

$$\|\mathbf{u} + \mathbf{v}\| \leq \|\mathbf{u}\| + \|\mathbf{v}\| \quad (\text{Triangle inequality for vectors})$$

$$d(\mathbf{u}, \mathbf{v}) \leq d(\mathbf{u}, \mathbf{w}) + d(\mathbf{v}, \mathbf{w}) \quad (\text{Triangle inequality for distances})$$

Proof: Consider $\|\mathbf{u} + \mathbf{v}\|^2$.

Orthogonality

Definition 5

Two vectors \mathbf{u}, \mathbf{v} are called **orthogonal** if $\langle \mathbf{u}, \mathbf{v} \rangle = 0$.

Question: What is the angle between any two orthogonal vectors?

Check for orthogonality

Example 6

Determine whether \mathbf{u} and \mathbf{v} are orthogonal or not, with the standard inner product.

① $\mathbf{u} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$ and $\mathbf{v} = \begin{pmatrix} 4 \\ 4 \\ -4 \end{pmatrix}$.

② $\mathbf{u} = \begin{pmatrix} 0 \\ 1 \\ 2 \\ 3 \end{pmatrix}$ and $\mathbf{v} = \begin{pmatrix} 5 \\ 2 \\ -3 \\ -4 \end{pmatrix}$.

③ $f = x$, $g = X^2$ on $[-1, 1]$.

④ $U = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$, $V = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$

Pythagorean Theorem

Theorem 7

If \mathbf{u}, \mathbf{v} are orthogonal vectors, then

$$\|\mathbf{u} + \mathbf{v}\|^2 = \|\mathbf{u}\|^2 + \|\mathbf{v}\|^2$$

Proof: Consider $\|\mathbf{u} + \mathbf{v}\|^2 = \langle \mathbf{u} + \mathbf{v}, \mathbf{u} + \mathbf{v} \rangle$.

Dr. Abdulla Eid

Example 8

Show that if \mathbf{u}, \mathbf{v} are orthogonal unit vectors in V , then $\|\mathbf{u} - \mathbf{v}\| = \sqrt{2}$.

Dr. Abdulla Eid

Orthogonal Complement

Definition 9

If W is a subspace of an inner vector space V , then the set of all vectors in V that are orthogonal to every vector in W is called the **orthogonal complement** of W and is denoted by W^\perp .

$$W^\perp := \{\hat{w} \in V \mid \langle \hat{w}, w \rangle = 0, \text{ for all } w \in W\}$$

Theorem 10

- 1 W^\perp is a subspace of W .
- 2 $W \cap W^\perp = \{\mathbf{0}\}$
- 3 $(W^\perp)^\perp$

Row space and null space are orthogonal

Example 11

Let $W = \text{span}\{\mathbf{w}_1, \mathbf{w}_2, \mathbf{w}_3\}$ where,

$$\mathbf{w}_1 = \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix}, \mathbf{w}_2 = \begin{pmatrix} -1 \\ -4 \\ 2 \end{pmatrix}, \mathbf{w}_3 = \begin{pmatrix} 4 \\ -5 \\ 13 \end{pmatrix}$$

Dr. Abdulla

Row space and null space are orthogonal

Example 12

Let $W = \text{span}\{\mathbf{w}_1, \mathbf{w}_2, \mathbf{w}_3\}$ where,

$$\mathbf{w}_1 = \begin{pmatrix} 3 \\ 0 \\ 1 \\ -2 \end{pmatrix}, \mathbf{w}_2 = \begin{pmatrix} -1 \\ -2 \\ -2 \\ 1 \end{pmatrix}, \mathbf{w}_3 = \begin{pmatrix} 4 \\ 2 \\ 3 \\ -3 \end{pmatrix}$$

Dr. Abdulla