

Section 1.5

Inverse of a matrix

Dr. Abdulla Eid

College of Science

MATHS 211: Linear Algebra

Goal:

- 1 To define the inverse of a matrix.
- 2 To find the inverse of a matrix.
- 3 To solve linear system using the inverse of a matrix.
- 4 To explore some of the properties of the inverse of matrices.

1 - Definition of the inverse of a matrix

Recall:

- If a is a real number, then the **additive inverse** of a is $-a$, such that

$$a + (-a) = 0 \text{ and } (-a) + a = 0$$

- If a is a **nonzero** real number, then the **multiplicative inverse** of a is $\frac{1}{a}$, such that

$$a \cdot \frac{1}{a} = 1 \text{ and } \frac{1}{a} \cdot a = 1$$

- If f is a function **passing the horizontal line test**, then the inverse of f is f^{-1} such that

$$(f \circ f^{-1})(x) = x \text{ and } (f^{-1} \circ f)(x) = x$$

Definition 1

Let A be an $n \times n$ -matrix. The **inverse matrix** (if it exists) of A is another matrix A^{-1} such that

$$A \cdot A^{-1} = I_n \text{ and } A^{-1} \cdot A = I_n$$

2 - How to find the inverse of a matrix

We write

$$(A|I_n) \tag{1}$$

and then we reduce (1) to get,

$$(I_n|A^{-1})$$

Note: If we can't reduce (1), then the matrix has **no** inverse.

Example 2

Find A^{-1} for

$$A = \begin{pmatrix} 3 & 1 \\ 4 & 1 \end{pmatrix}$$

Solution:

$$\begin{aligned} & \left(\begin{array}{cc|cc} 3 & 1 & 1 & 0 \\ 4 & 1 & 0 & 1 \end{array} \right), & R_1 & \rightarrow \frac{1}{3}R_1 \\ & \left(\begin{array}{cc|cc} 1 & \frac{1}{3} & \frac{1}{3} & 0 \\ 4 & 1 & 0 & 1 \end{array} \right), & R_2 & \rightarrow R_2 - 4R_1 \\ & \left(\begin{array}{cc|cc} 1 & \frac{1}{3} & \frac{1}{3} & 0 \\ 4 - 4(1) & 1 - 4(\frac{1}{3}) & 0 - 4(\frac{1}{3}) & 1 - 4(0) \end{array} \right), \\ & \left(\begin{array}{cc|cc} 1 & \frac{1}{3} & \frac{1}{3} & 0 \\ 0 & \frac{-1}{3} & \frac{-4}{3} & 1 \end{array} \right), & R_2 & \rightarrow \frac{1}{\frac{-1}{3}}R_2 \\ & \left(\begin{array}{cc|cc} 1 & \frac{1}{3} & \frac{1}{3} & 0 \\ 0 & 1 & 4 & -3 \end{array} \right), & R_1 & \rightarrow R_1 - \frac{1}{3}R_2 \end{aligned}$$

$$\begin{pmatrix} 1 & \frac{1}{3} & | & \frac{1}{3} & 0 \\ 0 & 1 & | & 4 & -3 \end{pmatrix}, \quad R_1 \rightarrow R_1 - \frac{1}{3}R_2$$

$$\begin{pmatrix} 1 & \frac{1}{3} - \frac{1}{3} & | & \frac{1}{3} - \frac{1}{3}(4) & 0 - \frac{1}{3}(-3) \\ 0 & 1 & | & 4 & -3 \end{pmatrix},$$

$$\begin{pmatrix} 1 & 0 & | & -1 & 1 \\ 0 & 1 & | & 4 & -3 \end{pmatrix}$$

Thus,

$$A^{-1} = \begin{pmatrix} -1 & 1 \\ 4 & -3 \end{pmatrix}$$

To check our answer, we must show that $AA^{-1} = I_n$ and $A^{-1}A = I_n$. We have that $AA^{-1} =$

$$\begin{pmatrix} 3 & 1 \\ 4 & 1 \end{pmatrix} \begin{pmatrix} -1 & 1 \\ 4 & -3 \end{pmatrix} \\ = \begin{pmatrix} 3(-1) + 1(4) & 3(1) + 1(-1) \\ 4(-1) + 1(4) & 4(1) + 1(-3) \end{pmatrix} \\ = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = I_2$$

To check our answer, we must show that $AA^{-1} = I_n$ and $A^{-1}A = I_n$. We have that $A^{-1}A =$

$$\begin{pmatrix} -1 & 1 \\ 4 & -3 \end{pmatrix} \begin{pmatrix} 3 & 1 \\ 4 & 1 \end{pmatrix} \\ = \begin{pmatrix} -1(3) + 1(4) & -1(1) + 1(1) \\ 4(3) + -3(4) & 4(1) + -3(1) \end{pmatrix} \\ = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = I_2$$

Example 3

Find A^{-1} for

$$A = \begin{pmatrix} 1 & 1 \\ 3 & 3 \end{pmatrix}$$

Solution:

$$\begin{pmatrix} 1 & 1 & | & 1 & 0 \\ 3 & 3 & | & 0 & 1 \end{pmatrix}, \quad R_2 \rightarrow R_2 - 3R_1$$
$$\begin{pmatrix} 1 & 1 & | & 1 & 0 \\ 3 - 3(1) & 3 - 3(1) & | & 0 - 3(1) & 1 - 3(0) \end{pmatrix},$$
$$\begin{pmatrix} 1 & 1 & | & 1 & 0 \\ 0 & 0 & | & -3 & 1 \end{pmatrix},$$

Since we couldn't reduce the matrix above, then it has **no** inverse!

Exercise 4

Find the inverse of

$$A = \begin{pmatrix} 1 & 0 \\ a & 1 \end{pmatrix}, \quad a \in (-\infty, \infty)$$

Dr. Abdulla Eid

Example 5

Find A^{-1} for

$$A = \begin{pmatrix} 2 & 1 & 0 \\ 4 & -1 & 5 \\ 1 & -1 & 2 \end{pmatrix}$$

Solution:

$$\left(\begin{array}{ccc|ccc} 2 & 1 & 0 & 1 & 0 & 0 \\ 4 & -1 & 5 & 0 & 1 & 0 \\ 1 & -1 & 2 & 0 & 0 & 1 \end{array} \right),$$

$$R_1 \leftrightarrow R_3$$

$$\left(\begin{array}{ccc|ccc} 1 & -1 & 2 & 0 & 0 & 1 \\ 4 & -1 & 5 & 0 & 1 & 0 \\ 2 & 1 & 0 & 1 & 0 & 0 \end{array} \right),$$

$$R_2 \rightarrow R_2 - 4R_1, R_3 \rightarrow R_3 - 2R_1$$

$$\left(\begin{array}{ccc|ccc} 1 & -1 & 2 & 0 & 0 & 1 \\ 4 - 4(1) & -1 - 4(-1) & 5 - 4(2) & 0 - 4(0) & 1 - 4(0) & 0 - 4(1) \\ 2 - 2(1) & 1 - 2(-1) & 0 - 2(2) & 1 - 2(0) & 0 - 2(0) & 0 - 2(1) \end{array} \right),$$

$$\left(\begin{array}{ccc|ccc} 1 & -1 & 2 & 0 & 0 & 1 \\ 0 & 3 & -3 & 0 & 1 & -4 \\ 0 & 3 & -4 & 1 & 0 & -2 \end{array} \right), \quad R_2 \rightarrow \frac{1}{3}R_2$$

$$\left(\begin{array}{ccc|ccc} 1 & -1 & 2 & 0 & 0 & 1 \\ 0 & 1 & -1 & 0 & \frac{1}{3} & \frac{-4}{3} \\ 0 & 3 & -4 & 1 & 0 & -2 \end{array} \right), \quad R_3 \rightarrow R_3 - 3R_2, R_1 \rightarrow R_1 + R_2$$

$$\left(\begin{array}{ccc|ccc} 1+0 & -1+1 & 2+(-1) & 0+0 & 0+\frac{1}{3} & 1+\frac{-4}{3} \\ 0 & 1 & -1 & 0 & \frac{1}{3} & \frac{-4}{3} \\ 0 & 3-3(1) & -4-3(-1) & 1-3(0) & 0-3(\frac{1}{3}) & -2-3(\frac{-4}{3}) \end{array} \right),$$

$$\left(\begin{array}{ccc|ccc} 1 & 0 & 1 & 0 & \frac{1}{3} & \frac{-1}{3} \\ 0 & 1 & -1 & 0 & \frac{1}{3} & \frac{-4}{3} \\ 0 & 0 & -1 & 1 & -1 & 2 \end{array} \right), \quad R_3 \rightarrow -R_3$$

$$\left(\begin{array}{ccc|cc} 1 & 0 & 1 & 0 & \frac{1}{3} & -\frac{1}{3} \\ 0 & 1 & -1 & 0 & \frac{1}{3} & -\frac{4}{3} \\ 0 & 0 & 1 & -1 & 1 & -2 \end{array} \right), \quad R_2 \rightarrow R_2 + R_3, R_1 \rightarrow R_1 - R_3$$

$$\left(\begin{array}{ccc|cc} 1-0 & 0-0 & 1-1 & 0-(-1) & \frac{1}{3}-1 & -\frac{1}{3}-(-2) \\ 0+0 & 1+0 & -1+1 & 0+(-1) & \frac{1}{3}+1 & -\frac{4}{3}+(-2) \\ 0 & 0 & 1 & -1 & 1 & -2 \end{array} \right),$$

$$\left(\begin{array}{ccc|cc} 1 & 0 & 0 & 1 & -\frac{2}{3} & \frac{5}{3} \\ 0 & 1 & 0 & -1 & \frac{4}{3} & -\frac{10}{3} \\ 0 & 0 & 1 & -1 & 1 & -2 \end{array} \right),$$

Thus,

$$A^{-1} = \begin{pmatrix} 1 & -\frac{2}{3} & \frac{5}{3} \\ -1 & \frac{4}{3} & -\frac{10}{3} \\ -1 & 1 & -2 \end{pmatrix}$$

To check our answer, we must show that $AA^{-1} = I_n$ and $A^{-1}A = I_n$. We have that $AA^{-1} =$

$$\begin{pmatrix} 1 & \frac{-2}{3} & \frac{5}{3} \\ -1 & \frac{3}{4} & \frac{-10}{3} \\ -1 & 1 & -2 \end{pmatrix}$$

$$\begin{pmatrix} 2 & 1 & 0 \\ 4 & -1 & 5 \\ 1 & -1 & 2 \end{pmatrix} \begin{pmatrix} 2(1) + 1(-1) + 0(-1) & 2(\frac{-2}{3}) + 1(\frac{3}{4}) + 0(1) & 2(\frac{5}{3}) + 1(\frac{-10}{3}) + 0(1) \\ 4(1) + (-1)(-1) + 5(-1) & 4(\frac{-2}{3}) + (-1)(\frac{3}{4}) + 5(1) & 4(\frac{5}{3}) + (-1)(\frac{-10}{3}) + 5(1) \\ 1(1) + (-1)(-1) + 2(-1) & 1(\frac{-2}{3}) + (-1)(\frac{3}{4}) + 2(1) & 1(\frac{5}{3}) + (-1)(\frac{-10}{3}) + 2(1) \end{pmatrix}$$

$$= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = I_3$$

To check our answer, we must show that $AA^{-1} = I_n$ and $A^{-1}A = I_n$. We have that $A^{-1}A =$

$$\begin{pmatrix} 1 & \frac{-2}{3} & \frac{5}{3} \\ -1 & \frac{4}{3} & \frac{-10}{3} \\ -1 & 1 & -2 \end{pmatrix} \begin{pmatrix} 2 & 1 & 0 \\ 4 & -1 & 5 \\ 1 & -1 & 2 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = I_3$$

3 - Solving Linear System using the inverse of a matrix

Example 6

Solve

$$3x + y = 2$$

$$4x + y = 3$$

Solution: This system can be written in a matrix multiplication form as

$$\begin{pmatrix} 3 & 1 \\ 4 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 2 \\ 3 \end{pmatrix}$$

$$A \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 2 \\ 3 \end{pmatrix}$$

$$A^{-1}A \begin{pmatrix} x \\ y \end{pmatrix} = A^{-1} \begin{pmatrix} 2 \\ 3 \end{pmatrix}$$

$$I_2 \begin{pmatrix} x \\ y \end{pmatrix} = A^{-1} \begin{pmatrix} 2 \\ 3 \end{pmatrix}$$

$$\begin{pmatrix} x \\ y \end{pmatrix} = A^{-1} \begin{pmatrix} 2 \\ 3 \end{pmatrix} = \begin{pmatrix} -1 & 1 \\ 4 & -3 \end{pmatrix} \begin{pmatrix} 2 \\ 3 \end{pmatrix}$$

$$\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1 \\ -1 \end{pmatrix}$$

Exercise 7

Solve the following system using the inverse matrix method.

$$2x - 3y = 9$$

$$4x + y = 1$$

Dr. Abdulla

Properties

- 1 The inverse is unique. We already have seen this!
- 2 (Socks-shoes property) $(AB)^{-1} = B^{-1}A^{-1}$.
- 3 $(A^{-1})^{-1} = I_n$. Why?
- 4 $A^n = AA \dots A$ and $A^{-n} = A^{-1}A^{-1} \dots A^{-1}$.
- 5 $(A^T)^{-1} = (A^{-1})^T$